

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) 21-08-2003		2. REPORT TYPE Technical Viewgraph Presentation		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Isolation of Boron and Carbon Atoms in Cryogenic Solids				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) C.W. Larson (AFRL/PRSP)				5d. PROJECT NUMBER 2303	
				5e. TASK NUMBER M2C8	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-PR-ED-VG-2003-212	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) AFRL-PR-ED-VG-2003-212	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES For presentation at the 9 th International Workshop of Combustion & Propulsion – Energetic Materials in La Spezia, Italy, taking place 14-19 September 2003.					
14. ABSTRACT					
20030929 082					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	A	14	Leilani Richardson
					19b. TELEPHONE NUMBER (include area code) (661) 275-5015

Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. Z39.18

Best Available Copy

Isolation of Boron and Carbon Atoms in Cryogenic Solids

**C. William Larson
Propulsion Directorate
Air Force Research Laboratory
Edwards AFB, CA 93524-7680**

**9th International Workshop on Combustion and Propulsion
NOVEL ENERGETIC MATERIALS AND APPLICATIONS
14-18 September 2003
Lerici, La Spezia, Italy**

Approved for public release; distribution unlimited.

Outline

Theoretical Isp of cryogenic solid propellants composed of the atoms, dimers and trimers of lightweight elements isolated in solid para hydrogen. Consequences of condensation.

Spectroscopic studies of Boron/Carbon clusters by matrix isolation spectroscopy.

Development of stable, hi-flux boron atom source for preparation of cryogenic solid HEDM (under auspices of Small Business Innovative Research (SBIR) program).


First optical spectrum of B_3 (under auspices of International Research Initiative of the Air Force Office of Scientific Research).

Video of exploding B/C and C HEDM.

Approved for public release; distribution unlimited.


Cryosolid Propellants Approach (Make)






- * Rapid vapor deposition of metal atom vapor and pre-cooled parahydrogen gas onto a liquid helium cooled substrate in vacuum.



Approved for public release; distribution unlimited.

Approved for public release; distribution unlimited.






Optimization of boron HEDM propellant combustion with liquid oxygen.

The propellant formulation is $H_{190}B_5$, or 5 equivalent mole percent boron atoms isolated in 95 mole percent solid parahydrogen. The four panels show the optimization for each of four levels of atom condensation: (1) B atoms, (2) B_2 molecules, (3) B_3 molecules, and (4) B_4 molecules. The I_{sp} and T_c were calculated for the Standard Rocket Condition: 1000 psi chamber pressure and expansion to sea level, which for LOX/LH₂ produces an I_{sp} of 389 s and a chamber temperature of 2984 K. The heats of formation for B_5H_{190} listed in each panel are derived from -2.20 kcal/mol for solid parahydrogen at 4.4 K, and 135.0 for B, 203.4 for B_2 , 192.8 for B_3 , and 225 kcal/mol for B_4 . The I_{sp} and T_c for no oxidizer are listed together with the optimum (maximum) I_{sp} obtainable for the specified O/F ratio (by mass) and the value of T_c . In all cases the chamber temperature with boron HEDM is very much less than the T_c of the LOX/LH₂ Standard Rocket, which produces $I_{sp} = 389$ s with $T_c = 2984$. The uncondensed boron HEDM I_{sp} of 474 s runs at 1832 K. With no oxidizer, the uncondensed boron HEDM rocket runs at 965 K and produces $I_{sp} = 402$ s.

Preparation



Annealing


a0 10 K	a3 32.5 K, 60 s	a6 40.0 K, 20 s
a1 27.5 K, 120 s	a4 35.0 K, 45 s	sublimation
a2 30.0 K, 90 s	a5 37.5 K, 20 s	rate ~ 1 $\mu\text{m}/\text{s}$

Precision matched pair of matrices


Green Matrix	$^{11}\text{B}/^{10}\text{B} = 80/20$	enhanced $^{11}\text{B}_j\text{C}_{n-j}$
Red Matrix	$^{11}\text{B}/^{10}\text{B} = 27/73$	enhanced $^{10}\text{B}_j\text{C}_{n-j}$

Approved for public release; distribution unlimited.

GOAL - 5% atoms in matrix




Approved for public release; distribution unlimited.



Approved for public release; distribution unlimited.

Approved for public release; distribution unlimited.









Figure 2. Annealing behaviors of B_xC_y species in matrix (a). Spectra labeled '1' were obtained from the originally deposited matrix, and spectra labeled '1' to '6' were obtained after successive annealings as detailed in the Fig. 1 caption. Absolute absorbance scales, $A_{\text{exp}} = \log(I_0/I)$, are offset to force coincidence at the peak maxima. Binary isotopomers of BC_4 and B_2C_4 are unresolved. The weaker of two bands of B_2C_3 ($\nu_{\text{asym}} = 1034 \text{ cm}^{-1}$, at 1830 cm^{-1}) is shown here. Spectral resolution is limited to $\sim 1 \text{ cm}^{-1}$ by matrix broadening.

Approved for public release; distribution unlimited.




Approved for public release; distribution unlimited.



Electronic absorption spectra recorded in a 6 K matrix after 4 hours of mass-selected co-deposition of B_3^- with neon. The bottom trace shows the $1E' - X^1A_1$ electronic transition of B_3^- overlapped by the $2^2E' - X^2A_1$ system of B_3 , produced from partial neutralization of the anions impinging on the matrix during deposition. The top trace reveals the $2^2E' - X^2A_1$ electronic transition of B_3 measured after exposure to UV radiation: Absorption belonging to the anion disappears.

M. Wyss, E. Riaplov, A. Batalov, J. P. Maier, T. Weber, W. Meyer, P. Rosmus, *J. Chem. Phys.* (2003, in press).
University of Basel, University of Kaiserslautern, Université de Marne la Vallée

Approved for public release; distribution unlimited.



Electronic absorption spectrum of the $1^2E' - X^2A_1'$ electronic transition of B_3 , recorded after 4 hours of mass-selected co-deposition with neon followed by UV irradiation of the 6 K matrix.

M. Wyss, E. Riaplov, A. Batalov, J. P. Maier, T. Weber, W. Meyer, P. Rosmus, *J. Chem. Phys.* (2003, in press).
University of Basel, University of Kaiserslautern, Université de Marne la Vallée

Approved for public release; distribution unlimited.

AFRL-PR-ED-TTR-2003-0030

AFRL-PR-ED-TTR-2003-0030

Advanced Rocket Propulsion Technologies Boron Vapor Source for HEDM

Paul C. Nordine

Containerless Research Inc.
906 University Place
Evanston IL 60201-3149

June 2003


SBIR Phase I Final Report

APPROVED FOR PUBLIC RELEASE. DISTRIBUTION UNLIMITED.



AIR FORCE RESEARCH LABORATORY
AIR FORCE MATERIEL COMMAND
EDWARDS AIR FORCE BASE CA 93524-7048

Approved for public release; distribution unlimited.



Approved for public release; distribution unlimited.

Conclusions

Large Isp improvements are produced by cryogenic solid propellants with atoms, dimers, trimers, and tetramers isolated in solid hydrogen, but condensation leads to loss of benefit.

5 mole percent B atoms produces Isp of 474 seconds compared to 389 s for LOX/sH₂. The HEDM combustion temperature is 1832 K, compared to 2984 K for LOX/sH₂.

Annealing kinetics of disappearance of C₃ and BC₂, and of appearance of B₂C, C₄, BC₃, B₂C₂, C₅, BC₄, and B₂C₃ unequivocally establishes the presence of atoms and dimers in the originally deposited matrix.

~80% or more of the initially deposited HEDM existed as atoms, dimers and trimers.

B₂C_n molecules are linear, with boron atoms attached to each end, and are immune to radical attack and condensation during annealing.

Theory predicts that a 12 kcal/mol barrier exists for B atom insertion into H₂, so isolation by co-condensation may be possible.

A stable, high-flux B-atom source has been developed under the Small Business Innovative Research Program capable of production of 100 mg of Boron HEDM in a few hours.

B₂ or B₃ may be the ultimate sinks (islands of stability) for atoms in the low temperature environment.

Studies of the spectroscopy and reactivity of B atoms and small clusters with hydrogen are underway at the University of Basel, supported by the Air Force Office of Scientific Research through the International Research Initiative program.

Approved for public release; distribution unlimited.

BACKUP CHARTS


Approved for public release; distribution unlimited.

Species	ΔH_f	ΔG_f	% LOX	$\Delta H_{f,2}$	% M	% LOX	% H_2
H_2 (g)	-22.1	—	—	336	0	20.6	79.4
H	52.1	107	19.0	1478	100.0	0	0
LiH	34.2	399	13.1	578	20.4	5.1	79.6
BeH	82.4	155	2.8	451	15.2	7.6	84.8
BH	109.3	410	3.8	568	28.3	0	71.7
CH	143.2	338	2.2	609	24.8	0	75.2
MgH	55.7	106	14.9	468	14.0	7.0	86.0
AlH	61.2	315	10.1	445	11.1	8.4	88.9
Li	38.1	407	12.2	451	19.9	5.0	80.1
Li_2	53.6	167	5.8	161	11.6	5.8	88.4
LiBe	109.8	182	3.9	572	15.0	7.6	85.0
LiB	159.6	434	5.0	527	29.0	0	71.0
LiC	159.9	435	1.3	529	30.0	0	70.0
LiMg	69.3	401	5.8	482	8.3	6.2	91.7
LiAl	97.7	448	5.0	458	7.3	7.4	92.7
Be	77.4	152	2.5	341	14.4	7.2	85.6
Be_2	153.1	325	5.0	545	7.8	7.8	92.2
BeAl	147.4	438	6.3	435	6.2	7.7	93.8
B	135.0	172	3.8	607	23.0	0	77.0
B_2	207.2	492	7.4	550	14.3	0	85.7
BC	201.6	182	3.7	542	14.2	0	85.8
C	171.3	469	0.0	645	20.0	0	80.0
C_2	199.3	462	0.0	533	15.3	0	84.7
CAI	174.5	458	3.8	464	6.8	5.1	93.2
N	113.0	414	15.0	551	34.2	0	65.8
Mg	35.2	398	16.8	417	13.8	7.1	86.2
Mg_2	68.8	408	8.9	416	7.4	7.6	92.6
Al	78.9	425	7.5	456	10.2	7.7	89.8
Al_2	125.1	445	7.5	445	5.6	8.4	94.4
Si	107.6	432	5.1	451	8.2	8.2	91.8
Ti	113.2	404	11.5	414	9.0	7.9	91.0

Conditions: Chamber Pressure = 1000 psi, Exhaust Pressure = 14.7 psi

Approved for public release; distribution unlimited.


Theoretical Infrared Intensities Linear C_n, DFT/B3LYP




Approved for public release; distribution unlimited.

TCNInte3.axg


Carbon cluster distributions




Approved for public release; distribution unlimited.



Survey spectra of precision matched matrices showing larger clusters $B_J C_{n-J}$, $n > 4$, $J = 0, 1, 2$ in original matrices and after three annealings.




Identification of 9 of the 16 isotopomers of linear BCCC in 5 matrices.




Approved for public release; distribution unlimited.

Identification of 7 isotopomers of the 10 isotopomers of BCCB in 5 matrices.



Approved for public release; distribution unlimited.


Four minimum energy geometries of B_2C_2 produce similar isotopomer fingerprints.
 Scale factor (S_F = measured frequency/theoretical frequency) of linear BCCB = 0.97250.



Approved for public release; distribution unlimited.

Figure 2 Phase 1 boron vapor source apparatus.

Approved for public release; distribution unlimited.



ITEM	QTY	DESCRIPTION
1	1	PHASE 1 CHAMBER
2	1	1/4" CP 316 ST. NUT
3	1	1/4" CP 316 ST. NUT
4	1	1/4" CP 316 ST. NUT
5	1	1/4" CP 316 ST. NUT
6	1	1/4" CP 316 ST. NUT
7	1	1/4" CP 316 ST. NUT
8	1	1/4" CP 316 ST. NUT
9	1	1/4" CP 316 ST. NUT
10	1	1/4" CP 316 ST. NUT
11	1	1/4" CP 316 ST. NUT
12	1	1/4" CP 316 ST. NUT
13	1	1/4" CP 316 ST. NUT
14	1	1/4" CP 316 ST. NUT
15	1	1/4" CP 316 ST. NUT
16	1	1/4" CP 316 ST. NUT
17	1	1/4" CP 316 ST. NUT
18	1	1/4" CP 316 ST. NUT
19	1	1/4" CP 316 ST. NUT
20	1	1/4" CP 316 ST. NUT
21	1	1/4" CP 316 ST. NUT
22	1	1/4" CP 316 ST. NUT
23	1	1/4" CP 316 ST. NUT
24	1	1/4" CP 316 ST. NUT
25	1	1/4" CP 316 ST. NUT
26	1	1/4" CP 316 ST. NUT
27	1	1/4" CP 316 ST. NUT
28	1	1/4" CP 316 ST. NUT
29	1	1/4" CP 316 ST. NUT
30	1	1/4" CP 316 ST. NUT
31	1	1/4" CP 316 ST. NUT
32	1	1/4" CP 316 ST. NUT
33	1	1/4" CP 316 ST. NUT
34	1	1/4" CP 316 ST. NUT
35	1	1/4" CP 316 ST. NUT
36	1	1/4" CP 316 ST. NUT
37	1	1/4" CP 316 ST. NUT
38	1	1/4" CP 316 ST. NUT
39	1	1/4" CP 316 ST. NUT
40	1	1/4" CP 316 ST. NUT
41	1	1/4" CP 316 ST. NUT
42	1	1/4" CP 316 ST. NUT
43	1	1/4" CP 316 ST. NUT
44	1	1/4" CP 316 ST. NUT
45	1	1/4" CP 316 ST. NUT
46	1	1/4" CP 316 ST. NUT
47	1	1/4" CP 316 ST. NUT
48	1	1/4" CP 316 ST. NUT
49	1	1/4" CP 316 ST. NUT
50	1	1/4" CP 316 ST. NUT
51	1	1/4" CP 316 ST. NUT
52	1	1/4" CP 316 ST. NUT
53	1	1/4" CP 316 ST. NUT
54	1	1/4" CP 316 ST. NUT
55	1	1/4" CP 316 ST. NUT
56	1	1/4" CP 316 ST. NUT
57	1	1/4" CP 316 ST. NUT
58	1	1/4" CP 316 ST. NUT
59	1	1/4" CP 316 ST. NUT
60	1	1/4" CP 316 ST. NUT
61	1	1/4" CP 316 ST. NUT
62	1	1/4" CP 316 ST. NUT
63	1	1/4" CP 316 ST. NUT
64	1	1/4" CP 316 ST. NUT
65	1	1/4" CP 316 ST. NUT
66	1	1/4" CP 316 ST. NUT
67	1	1/4" CP 316 ST. NUT
68	1	1/4" CP 316 ST. NUT
69	1	1/4" CP 316 ST. NUT
70	1	1/4" CP 316 ST. NUT
71	1	1/4" CP 316 ST. NUT
72	1	1/4" CP 316 ST. NUT
73	1	1/4" CP 316 ST. NUT
74	1	1/4" CP 316 ST. NUT
75	1	1/4" CP 316 ST. NUT
76	1	1/4" CP 316 ST. NUT
77	1	1/4" CP 316 ST. NUT
78	1	1/4" CP 316 ST. NUT
79	1	1/4" CP 316 ST. NUT
80	1	1/4" CP 316 ST. NUT
81	1	1/4" CP 316 ST. NUT
82	1	1/4" CP 316 ST. NUT
83	1	1/4" CP 316 ST. NUT
84	1	1/4" CP 316 ST. NUT
85	1	1/4" CP 316 ST. NUT
86	1	1/4" CP 316 ST. NUT
87	1	1/4" CP 316 ST. NUT
88	1	1/4" CP 316 ST. NUT
89	1	1/4" CP 316 ST. NUT
90	1	1/4" CP 316 ST. NUT
91	1	1/4" CP 316 ST. NUT
92	1	1/4" CP 316 ST. NUT
93	1	1/4" CP 316 ST. NUT
94	1	1/4" CP 316 ST. NUT
95	1	1/4" CP 316 ST. NUT
96	1	1/4" CP 316 ST. NUT
97	1	1/4" CP 316 ST. NUT
98	1	1/4" CP 316 ST. NUT
99	1	1/4" CP 316 ST. NUT
100	1	1/4" CP 316 ST. NUT